




Forestry Commission
England

South East England

Battle Great Wood

Design Concept

Illustrates the main features and broad character of the woodland in 30 years time.



Produced by the Planning Team 14th February 2008

This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.


N

1:10,000



Forestry Commission
England

South East England Beckley and Bixley Woods



Produced by the Planning Team 14th February 2008

This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N

1:12,000




Forestry Commission
England

South East England Bedgebury Forest

Design Concept


Illustrates the main features and broad character of the woodland in 30 years time.



Produced by the Planning Team 21st February 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:12,000



South East England Burwash (1)

Design Concept

Illustrates the main features and broad character of the woodland in 30 years time.

Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.

Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.

Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.

Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.

Wooded heath (20-70% tree canopy cover)

Predominantly conifer woodland retained beyond economic maturity for habitat diversity and landscape interest.

Open space

Management area

Watercourse


Open water

Produced by the Planning Team 19th February 2008

This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N

1:10,000



Forestry Commission
England

South East England
Burwash (2)

Design Concept

Illustrates the main features and broad character of the woodland in 30 years time.

Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.

Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.

Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.

Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.

Wooded heath (20-70% tree canopy cover)

Predominantly conifer woodland retained beyond economic maturity for habitat diversity and landscape interest.

Open space

Management area

Open water

— Watercourse

Produced by the Planning Team

19th February 2003

This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission 100025498, 2007.

N
1:15,000