










-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse

APPROVED BY (FC):

DATE:

Produced by the Planning Team 28th Feb 2008


This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.










N
1:10,000

South East England

Beckley and Bixley Woods

Future Habitats Map



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):

DATE:










Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:11,000



South East England Bedgebury Forest Future Habitats Map

-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Forest road
-  Compartments
-  Forest access tracks
-  Sub-compartments
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 20th March 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office © Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:11,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:15,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

APPROVED BY (NE):

DATE:

APPROVED BY (FC):

DATE:

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse

Produced by the Planning Team 28th Feb 2008


This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.










N
1:15,000

South East England

Harrison's Rocks (Birchden Wood)

Future Habitats Map



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):

DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:11,000




APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:12,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Outside of the SSSI, shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

APPROVED BY (NE):

DATE:

APPROVED BY (FC):


DATE:










-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse


APPROVED BY (FC):










DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000



-  Predominantly native woodland with a small element of conifer (<20%). Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Native broadleaf woodland (Natural Reserve).
-  Mixed woodland with native trees regenerating amongst and gradually replacing conifer woodland. Small patches of open space support early successional vegetation during the establishment of young native trees. Shifting areas of wooded heath support 20-70% tree canopy cover and are located in close proximity to the permanent network of environmental corridors.
-  Predominantly conifer woodland.
-  Sweet chestnut coppiced on a cycle provides cyclical open space with early successional vegetation during the regrowth phase of the coppice stools.
-  Permanent network of open habitat supports heather and associated species. These environmental corridors measure 10-40 metres in width and are integrated with patches of wooded heath. They also link up with patches of open habitat on both Forest Enterprise and neighbouring land.
-  Wooded heath (20-70% tree canopy cover)
-  Conifer specimens grown in mixture with open habitat in Bedgebury National Pinetum.
-  Buildings/car parks facilitating public access.

-  Open water
-  Compartments
-  Sub-compartments
-  Forest road
-  Forest access tracks
-  Watercourse

APPROVED BY (FC):


DATE:

Produced by the Planning Team 28th Feb 2008

This map is based upon Ordnance Survey material with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationary Office ©Crown Copyright. Unauthorised reproduction infringes Crown Copyright and may lead to prosecution or civil proceedings. Forestry Commission, 100025498, 2007.

N
1:10,000

Bedgebury Pinetum – Education and Conservation Plantings





Forest Enterprise
Bedgebury Office, Park Lane, Goudhurst,
Kent, TN17 2SL
www.forestry.gov.uk