

# Tamar Valley Forest Plan 2016 - 2026 West England Forest District



Forestry Commission woodlands have been certified in accordance with the rules of the Forest Stewardship Council.

The mark of responsible forestry PEFC/18-40-1001 Promoting Sustainable Forest Management www.pefc.org

**Declaration by FC as an Operator.** 

All timber arising from the Forest Enterprise estate represents a negligible risk under EUTR (No 995/210)

Ben Robinson FCE File Ref: OP10/83 OLD Ref: PE74 & 74/1



### **List of Contents**

#### **PART 1 – Description, summary & objectives**

| Application for Forest Plan Approval | 2 |
|--------------------------------------|---|
| Contents                             | З |
| Location                             | 4 |
| Summary                              | 5 |
| Management objectives                | 6 |
| Meeting Objectives                   | 7 |

#### **PART 2 – Character, analysis & concept**

| Landscape Character | 8  |
|---------------------|----|
| Designations        | 9  |
| Analysis & Concept  | 10 |

#### **PART 3 – Composition and future management**

| Woodland Composition | 11 |
|----------------------|----|
| Naturalness on PAWS  | 12 |
| PAWs Management      | 13 |
| Broadleaf Management | 14 |

### PART 4 – Thinning, felling and future composition

| Silviculture                             | 15 |
|--|----|
| Alternatives to Clearfell                | 16 |
| Management Prescriptions 2016-2046       | 17 |
| Indicative Future Species, 2026 and 2046 | 18 |

#### **PART 5 – Conservation**, heritage and recreation

| Conservation Habitats         | 19 |
|-------------------------------|----|
| Conservation features         | 20 |
| Heritage features             | 21 |
| Recreation, Tenure and Access | 22 |

| <b>APPENDIX</b> | 1:         | Physical | envi |
|-----------------|------------|----------|------|
| AFFLINDIA       | - <b>-</b> | FILYSICA |      |

| Geology            |
|--------------------|
| Soils              |
| Landform           |
| Landscape Analysis |

#### **APPENDIX 2: Management considerations**

| Option Testing     |  |  |
|--------------------|--|--|
| Stock data – 2016  |  |  |
| Pests and Diseases |  |  |

#### **APPENDIX 3: Supporting Information**

Glossary of Terms References

#### **APPENDIX 4: Consultation**

**Consultation Record** 

#### ironment

| 23  |    |
|-----|----|
| 23  |    |
| 24  |    |
| 25- | 26 |

| 27 |
|----|
| 28 |
| 29 |
|    |

| 30 | -31 |
|----|-----|
| 32 |     |

33



© Crown copyright and database right [2016] Ordnance Survey [100021242]

0 0.325 0.65 1.3 1.95 2.6 Miles

#### Tamar Valley Forest Plan 2016 - 2026 Page 4





## Location

The Tamar Valley Forest Plan area lies within West Devon District and is a signification feature of the Tamar Valley Area of Outstanding Natural Beauty. The Plan area is made up of the two distinct forest blocks of **Denham** and **Birch Wood** totalling 138ha. The forests are close to the settlements of Tavistock and Bere Alston.

| Forest Name | Area (ha) | % of Plan<br>Area |
|-------------|-----------|-------------------|
| Denham      | 98.8      | 72                |
| Birch Wood  | 39.2      | 28                |

The Plan area sits within a wooded valley landscape and provides both a visual feature and recreational attraction for the surrounding area. Both woodlands border the River Tavy which is a major tributary of the River Tamar.

The majority of the land is at 5-125 metres above sea level (asl) and is undulating to very steep in places. The climate is warm and fairly moist with an average annual rainfall of 1000— 1400mm, a soil moisture deficit of around 150mm, and an accumulated temperature over 5°C of 2000°C.

The soils are primarily upland brown earths (1u) underlaid with shallow rock within the River Tamar valley.

## Summary

### About

The Tamar Valley Forest Plan area is made up of two separate forest blocks totalling 138 hectares in Devon. The forests lie within the Tamar Valley Area of Outstanding Natural Beauty. As individual forest blocks set within the intimate wooded valley landscape they have very high natural and landscape diversity and value.

The forests managed as part of the public forest estate are Denham (99ha), close to Bere Alston and Buckland Abbey and Birch Wood (39ha) in which is 2 miles south of Tavistock.

The public forest here is a predominantly conifer on ancient woodland (PAWS) having been planted to address the national timber shortage of the early Twentieth Century. The area is known to produce high quality Douglas fir which makes up the majority of the trees here. Areas of remnant ancient semi-natural woodland do remain and are made up of oak, ash and sweet chestnut. Most of the areas are actively managed to provide timber for local and national businesses, and to improve the quality of the remaining tree crop.

The Plan area contains numerous archaeological features, notably mines, which originate from the rich iron and silver mining for which the area was renowned.

The Plan area is a rich for ecology and includes Priority Lowland Mixed Deciduous Woodland which is habitat in part for dormice.

No Public Rights of Way cross the Plan area but Birch Wood is Open Access, with Denham being de facto Open Access due to the nature of the landholding. The majority of recreation usage is made up of walkers with some limited amount of usage by horse riders and mountain bike riders.

### **Objectives**

The core aim of the plan is to produce woodlands with increased conservation and landscape benefits whilst maintaining a viable timber output. The long term aims of management here are to continue the process of restoring Ancient Woodland while increasing resilience to climate, pest and disease risks, and to develop the forest for people. The social, economic and environmental objectives of management here are to:

- The continued production of sustainable and marketable woodland products.
- To conserve, maintain and enhance cultural and heritage assets.
- The provision and maintenance of recreation facilities.
- Protect and enhance woodland and open habitats and their associated species.
- To protect enhance and restore areas of ancient woodland in line with the 2005 'Keepers of Time' policy.
- The delivery of well-designed proposals in keeping with AONB and local landscape character.

### What we'll do

The current plan outlines management proposals including felling and restocking over several decades, with felling licence approval for operations up until 2026.

Areas identified as PAWS will be managed as mixed woodland to maximise their productive potential, with the aim of a gradual return to native woodland. Prescribed proactive management of the broadleaf areas w

The Plan makes provision to ensure proposals are in keeping with the neighbouring intimate wooded landscape.

Implementation and maintenance of an environmental corridor system will continue to increase diversity of habitat and internal landscaping.

The planned areas of clearfelling, restocking and permanent open space creation during the ten years to 2026 are summarised in the chart below.

| HECTARES                | Conifers | Broadleaves | Open space |
|-------------------------|----------|-------------|------------|
| Clearfelling            | -        | 1.8         | -          |
| Restocking/Regeneration | -        | 1.8         | -          |

In addition to these defined operations, ongoing thinning and selective felling of both conifers and broadleaves will be carried out in the plan area at five to ten year intervals.

The proportions of conifer and broadleaved woodland and open space at the beginning of the plan period are shown in the bar chart. The increase in native broadleaves expected within the plan period and over time is indicated in the middle and right hand columns of the chart.

|   | 100 |   |
|---|-----|---|
|   | 90  |   |
|   | 80  |   |
|   | 70  |   |
|   | 60  |   |
| % | 50  |   |
|   | 40  | _ |
|   | 30  |   |
|   | 20  |   |
|   | 10  |   |
|   | 0   |   |







### **Management Objectives**

### WEST ENGLAND FOREST DISTRICT

**PROTECTING AND EXPANDING ENGLANDS FORESTS** AND WOODLANDS AND INCREASING THEIR VALUE TO SOCIETY AND THE ENVIRONMENT.

The objectives of this Plan will, in part, deliver the West England Forest District Strategic Plan (2013a) and the national Strategic Plan for the Public Forest Estate in England (2013b).

Sustainable management of the woodland will be to the standards required to maintain FSC and PEFC accreditation as stipulated in UKFS and UKWAS and therefore must deliver economic, environmental and social objectives.

following page.



responsible forestry

**Declaration by FC as an Operator.** 

All timber arising from the Forest Enterprise estate represents a negligible risk under EUTR (No 995/210)

The continued production of sustainable and marketable woodland products.

The provision and maintenance of recreation facilities.

**Deliver well-designed** forests in keeping with the AONB and local landscape character.

> To protect enhance and restore areas of ancient woodland in line with the 2005 'Keepers of Time' policy.

To conserve,

maintain and

cultural and

heritage assets.

enhance

Protect and enhance woodland and open habitats and their associated species.

Vature

#### **Tamar Valley Forest Plan** 2016 - 2026 Page 6





The meeting and monitoring of these objectives is outlined on the

Forestry Commission woodlands have been certified in accordance with the rules of the Forest Stewardship Council.



## **Meeting Objectives**

| Objective  | Meeting Objective  |  |
|--|--|--|
| The continued production of sustainable and marketable woodland products.                                      | The majority of the Plan area will remain productive through<br>thinning yield.<br>Minimal clearfell timber production will occur from the<br>conifers.  | Comparison of total<br>annual average) wit<br>(FP) five and ten-ye<br>Pre-thinning survey  |
| To protect enhance and restore areas<br>of ancient woodland in line with the<br>2005 `Keepers of Time' policy. | Targeted felling of conifer crops and suppression of non-<br>native regeneration to aid natural native regeneration and na-<br>tive species replanting   | Analysis and compar<br>through the Forest F  |
| To conserve, maintain and enhance cultural and heritage assets.  | Liaise with Devon Archaeology Service prior to commencement of works in proximity to heritage assets. Where appropriate limit shrub encroachment on features.  | Operational site plan<br>operations will help<br>Feature condition m<br>records updated.   |
| Deliver well-designed forests in<br>keeping with the AONB and local<br>landscape character.                    | Implementation of proposals will soften and better integrate the woodland with the surrounding landscape   | Fixed point photogra   |
| Protect and enhance woodland and open habitats and their associated species.                                   | <ul> <li>Felling together with a delayed restock program will continue to diversify stand and age structure.</li> <li>Operational site planning should highlight opportunities where conservation benefits can be delivered.</li> <li>Appropriate reinstatement works will be carried out once operations have been concluded.</li> <li>Creation of &gt;10% transitory and permanent open space</li> </ul> | Monitored via Revie<br>updated sightings.<br>Analysis and compa<br>the Forest Plan rev<br>Operational site pla<br>operations will help |
| The provision and maintenance of recreation facilities.  | Management of existing car parking and trail facilities will be<br>maintained by the Beat team.<br>Visitor numbers will be maintained.<br>Road and ride corridor and car park aesthetics enhanced and<br>maintained.   | Beat team will monit<br>car park, trails and s   |

#### Tamar Valley Forest Plan 2016 - 2026 Page 7



ediende have en rentitien in conduirce with the co of the Forcet wandship Council.



### Monitoring

production forecast yield (1,000m<sup>3</sup> th actual production at the Forest Plan ear review.

followed by thinning control sheets.

rison of SCDB `naturalness' scores Plan review process.

nning of harvesting and restocking monitor the effect of management.

nonitored through Review process and

aphy analysis at Forest Plan review stage

ew process, through local records and

arison of SCDB open space 10% through view process.

anning of harvesting and restocking p monitor the effect of management

itor usage and ensure the maintenance of signage.

### 151 South Devon National Character Assessment Profile Source: Natural England (2012)

At its core, South Devon is a fertile, agricultural landscape, with smooth, rounded hills separated by deep, wooded valleys; a patchwork landscape of arable and improved pasture. The Tamar Valley has some distinctive features relating to the topography, land use and underlying geology. Principally the granite ridge that runs across the Tamar Valley results in a gorge-like landscape, with thickly wooded sides and rocky outcrops.

The area contains 12,374 ha (10 per cent of NCA), including 3,446 ha of ancient woodland. There are important ancient woodland sites on valley and ria sides and substantial areas of ancient woodland are concentrated in the Dart and Erme valleys. Generally oak or ash dominate, while alder and hazel occur in damper areas. Sweet chestnut, beech and sycamore are also common.



### **Tamar Valley AONB**

Tamar Valley AONB is a rare valley and water landscape based around an unspoilt estuary, a unique wildlife resource and a landscape of distinctive heritage.

The AONB contains extensive areas of woodland, particularly on the steep valley sides. This is split between large tracts of well-managed coniferous forest and equally large areas of broadleaved woodland, much of it ancient semi-natural. Within this are nationally important species, including the heath fritillary butterfly, and regionally

endangered species like the nightjar.

The vision of the AONB is to ensure that by working together the Tamar Valley AONB is managed in a way that conserves and enhances its unique cultural identity and outstanding natural beauty for the benefit of all the people who live, work and visit the area, now and in the future.



#### CHARACTER DESCRIPTION—River Tavy Middle Valley Source: West Devon District Council (2008)

This is a landscape of contrasts, with ridges of higher land separated by the winding wooded valley of the River Tavy. Extensive dark mixed plantation covers its valley sides, creating a sense of isolation and secrecy. Watermeadows on the valley floors and pockets of broadleaved woodland add variations in colour and texture. Streams glide between craggy wooded banks covered in mosses and ferns, tumble over boulders and rocks, and flow under ancient bridges. Sunken lanes twist their way up the valley sides towards an open landscape of pastoral fields, with sudden long views across to Dartmoor or the River Tamar. Buckland Abbey is nestled into the valley side, surrounded by parkland. The rich and diverse landscape has a strong sense of history, with prehistoric hillforts, farms, estates, villages and industrial remains all contributing to its sense of time-depth.

This area is separated from the Tamar valley to the west by a narrow plateau which runs from Bere Alston to Lamerton. Its boundaries with adjacent landscape character areas are all fairly gradual. To the north are the Tamar Upland Fringe and Tavistock Dartmoor Fringes; to the east Central Dartmoor and Southern Dartmoor and Fringes; to the south Plymouth Farmland and the city of Plymouth and to the west the Lower Tamar and Tavy Valleys and the Middle Tamar Valley.

The area continues to provide much-needed recreational provision for tourists and the residents of Plymouth, but in a sensitive way with minimal impact on the landscape and biodiversity. The area's dramatic views and historic features are protected, understood and enhanced, with interpretation and public access where appropriate. Any development is carefully designed and sited to minimise its impact on the area's rural character. Sustainable agriculture and the maintenance of traditional landscape features are supported. Semi-natural habitats such as heathland, woodland and watermeadows are well-managed to maximise their biodiversity and resilience to climate change.



| Protect                               | Manage                          | Plan            |
|---------------------------------------|---------------------------------|-----------------|
| Protect the area's <b>long</b>        | Manage woodland (including      | Plan for the lo |
| riews across Dartmoor and             | through traditional techniques  | large conifer   |
| he Tavy and Tamar Valleys.            | such as coppicing) to           | through grad    |
| Protect the area's <b>scenic</b>      | maximise age diversity and to   | native broad    |
| <b>[uality</b> , and enhance its role | develop species composition     | retaining and   |
| is a setting for adjacent             | appropriate to the area;        | conifer planta  |
| protected landscapes.                 | encourage the reversion of      | (e.g. for mou   |
| Protect the area's <b>high</b>        | deciduous woodland on           | Plan to create  |
| evels of tranquillity,                | maturity, especially on ancient | natural habita  |
| esisting development which            | woodland sites                  | and woodland    |

would reduce them.

2

long-term restructuring of plantations in the Tavy valley lual restocking with a range of leaved species; consider promoting less prominent ations as recreational spaces untain biking).

e, extend and link semiats such as heathland, hedges ds, including through the reinstatement of lost field boundaries.

## Landscape Character

#### Tamar Valley Forest Plan 2016 - 2026 Page 8





copyright and database right [2016] Survey [100021242] © Crown c Ordnance 3

#### **Ancient Woodland**

The Plan area and local landscape is host to considerable areas of designated Ancient Woodland sites (AWS), both remnant ancient semi-natural woodland (ASNW) and plantation on ancient woodland sites (PAWS). Of the total Plan area, over 75% is designated Ancient Woodland (106ha) with almost the entirety of Denham designation. A large proportion of this Ancient Woodland (94ha), is PAWS having been planted with non-native conifer in the 1960s. The majority of ASNW areas are found in the valley sides or bottoms and are oak dominated.











### Area of Outstanding Natural Beauty

The Plan area sits within the Tamar Valley Area of Outstanding Natural Beauty (AONB). AONBs are designated under the Countryside & Rights of Way Act (2000), and along with National Parks they represent the finest examples of countryside in England and Wales. Designation seeks to conserve and enhance natural beauty; including flora and fauna, geology and landscape features, archaeology and architecture; whilst recognising the needs of the local community and economy.

The Tamar Valley AONB contains an estimated 3,668ha (20.2%) of woodland, significantly above the regional average. Woodland condition is variable but the Valley produces timber – especially Douglas Fir – of excellent quality and the potential productivity is certainly higher than at present. Conifer crops, providing they are planted sympathetically in the landscape, have a role in sustaining the economic viability of land-holdings. Sustainably managed coniferous and broadleaved forests will continue to provide fuel, timber, carbon storage, biodiversity and other ecosystem services.

Analysis: Numerous copper, silver and lead mines are found throughout the woodlands, and are a key feature of the Tamar Valley cultural heritage.

**Concept:** These features are fenced but visible to the public maintaining their cultural value whilst keeping people safe.

Mining areas are avoided during operations.



Analysis: Both woodlands sit within a wooded landscape. Surrounding woodlands, many of which are designated as Ancient Woodland, adjoin the Plan area.

Concept: Woodland operations will need to

be carefully planned and sensitively executed to ensure the woodlands are managed in keeping with the local landscape.



Analysis: An outbreak of Phytophthora ramorum was found in 2014 in European larch and Sweet chestnut—with symptomatic trees requiring removal.

**Concept:** The application of coppicing of Sweet chestnut and thus precipitating vigorous regrowth is not deemed appropriate in this area. Continued removal of symptomatic trees will occur.

Analysis: Quality productive Douglas fir is found throughout the woodlands and is a key feature of the Tamar Valley.

**Concept:** These stands will be managed through thinning as an alternative to clearfelling. Douglas fir regeneration will be accepted provided it does not adversely impact ancient woodland eatures or remnants.



Analysis: A small area of secondary woodland sits exposed in the western portion of Denham. The western sides of the woodland neighbours open agricultural pasture.

**Concept:** This area will be managed through CCF in light of the large ancient woodland designation in the rest of the woodland with an objective of mixed woodland.

> Analysis: Areas of remnant ancient semi-natural woodland are found scattered across the Plan area. These predominantly consist of oak with beech intrusion.

Concept: These areas will be managed to ensure that the features are preserved and enhanced in perpetuity

**Analysis:** The focus of recreational access is at the car park at the northern end of Denham. The area is in good condition but somewhat over shadow by the surrounding crops.

**Concept:** Steps will be taken to improve the setting of the Car Park and entrance to the



Hunter



**Analysis:** The River Tavy traverses and links the Plan area, providing steep valley sides and areas of wet woodland. The woodlands play a protection function in the river catchment, providing soil stabilisation and run off slow-down

Concept: These areas will be managed sensitively as riparian zones, where soil and nutrient stability are prioritised. The target will be to reach a maximum of 50% forest cover of site appropriate wet woodland species.



#### Tamar Valley Forest Plan 2016 - 2026 Page 10





### **Analysis & Concept**

Analysis: The topography of much of the Plan area is of a gradient, some areas are particularly steep in Denham

Concept: These areas will be managed sensitively through continuous cover

forestry to ensure soil stability is maintained.









Native & naturalized broadleaves

Open/other

Note: Beech, sycamore and sweet chestnut are considered to be not within their native range but are considered to be 'naturalised'

## **Woodland Composition**

The Plan area is conifer dominated with significant remnants of ancient semi-natural and native broadleaf features. Denham had contained some larch components until recently, but these have now been removed following *Phythophotora ramorum* infection and Statutory Plant Health Notice felling. The majority of conifer components are made up of quality Douglas fir (74ha) with Norway spruce and Western hemlock the major supplementary species. The broadleaf components are predominantly made up of oak, sweet chestnut and beech. Ash, birch and sycamore are evident as pioneer species within discrete areas of the Plan area.

The vast majority of conifer crops are around 50 years old with considerable levels of planting having occurred in the 1960s. As a result the woodlands are even-aged and thus economically and ecologically resilience poor. The thinning of conifer crops has ensured that understorey development is beginning to establish which in time will deliver a more structurally diverse woodland composition.

The broadleaf components of the Plan area comprise a mixture of ancient semi-natural oak, sweet chestnut and beech assemblages and younger plantings and regeneration. The majority of stands are even aged with understory development evident but not always establishing as a secondary crop. Where broadleaves feature within conifer crops these have been favoured and halo thinned to assist crown development.



#### Tamar Valley Forest Plan 2016 - 2026 Page 11



mmtation have at in rwith the forest p Council.





Class 1 - Semi-Natural Woodland & Restored Plantation Woodland (> 80% site native species)



Class 2 – Plantation Woodland (50 - 80% site native species)



Class 3 - Plantation Woodland (20 - 50% site native species)



Class 4 - Plantation Woodland (< 20% site native species)



Legend



Class 1 - > 80% Site Native Species Class 2 - 50-80% Site Native Species Class 3 - 20-50% Site Native Species Class 4 - <20% Site Native Species

Classes 2, 3 and 4 are classified as Plantations on Ancient Woodland Sites (PAWS). Areas of Semi-Natural Woodland and restored PAWS (Class 1 - > 80% site native species) are mostly found towards the bottom of valleys, in wetter riparian areas where the soils are richer.

The transformation of Classes 2, 3 and 4 AWS towards Class 1 is a key objective of this Plan and is in line with the Forestry Commission England, Keepers in Time Policy (Forestry Commission, 2005).

sites.

#### **Tamar Valley Forest Plan** 2016 - 2026 Page 12





### Naturalness on PAWS

Naturalness is the measure to show the percentage of site native tree species in a given area. This measure is used to record and monitor the condition and restoration of Ancient Woodland Sites previously planted with non-native species.

There has been gradual yet significant change in the naturalness of the AWS across the Plan area since 2005. The development of native species within stands has occurred in all three naturalness classes which is enabling the gradual restoration of PAWS



© Crown copyright and database right [2016] Ordnance Survey [100021242]

#### **Transition Zone**

The indicative proportion of native tree species is 50% or more of the crop. Removal of remaining conifer will be achieved through repeated thinning operations.

The establishment period to predominantly native woodland within this category is anticipated to be 20 – 30 years but is dependant on successful regeneration and establishment although maybe sooner depending on the level of conifer needing to be removed. Scattered individual conifers or small groups may remain.

#### **Preparation Zone**

Areas within this category contain less than 50% of native tree species but have a proportion greater than 20% of the crop and the area neighbours an area of significant native species

cover which can be utilised as a seed source. Enhancement of native content will continue through thinning of the conifer content.

These areas will be thinned heavily to release ancient woodland remnants and features and to encourage natural regeneration and intrusion in to the non-native crop.

The anticipated time scale for establishment of predominantly native species is expected be around 50 – 60 years or so, but could be as long as 70 - 80 depending on success of establishing the future crop.

#### **Non-native Zone**

The proportion of native tree species within a management area is less than 20% of the crop. Thinning in both these sub-categories should encourage crown development of broadleaf

components. Progress will be monitored and crops moved into either depending on development of stand structure and the response of natural regeneration.

#### **Clearfell Zone**

One clearfell will be used to convert PAWS in 15-20 years time. This is felling of Western hemlock and is required to ensure the integrity of the coupe which is predominantly secondary

woodland. This will be restocked with site suitable native species

### 0 0.05 0.1 0.2 0.3 0.4

#### Birch Wood



Restoration of Plantations on Ancient Woodland Sites (PAWS) has already begun and this continued restoration is going to take a considerable amount of time and resource because of the limited native remnants from which sites can regenerate.

Therefore a proactive yet realistic approach will be used to transform these sites over a period of time.

The aim of the transitional period to woodland containing 80% or more of native species should be to achieve:





#### Tamar Valley Forest Plan 2016 - 2026 Page 13





### **PAWS Management**

- a varied age structure with varying ratios of high canopy, secondary canopy and understory through out.
  - transition that ensures a minimum future content of 3 native species, with 4 to 5 species being the preferable target.
  - a minimal reliance on monocultures especially of birch, ash, hazel or oak. In practice this may involve either underplanting or group felling and planting within existing mid rotation broadleaf crops.
  - restoration of beech and sweet chestnut stands will not be prioritised as these species are to be naturalised and offer greater broadleaf diversity and therefore resilience.

If adequate regeneration is not evident in the 'Transition' and 'Preparation' zones after 10 years a reappraisal of the prescription will be needed.

### Legend

Building Block (native seed source)
Transition Zone
Prepartion Zone
Non-native Zone
Clearfell





### **Broadleaf Management**

### Legend



The majority of the Plan area consists of National Vegetation Classification (NVC) types W10 and W16 oak woodlands with some W8 ash woodlands along lower slopes and valley bottoms. These classifications give a good indication of the target future species for PAWS restoration and if sites were left to natural succession.

These sites will be managed on shelterwood systems whereby the new crop will be regenerated from selected seed trees, or through coppicing of oak, sweet chestnut and hazel stools where appropriate. In areas where the removal of Sweet chestnut symptomatic of *Phytophthora ramorum is required* the application of coppicing is not deemed appropriate in this area. Continued removal of symptomatic trees will occur. Light levels will be managed to minimise weed encroachment and grazing pressure from deer to limit regeneration predation following thinning operations.

Underplanting with species such as hazel, lime and hornbeam may be considered on the few ash dominated sites to ensure greater resilience to *Chalara fraxinea*. The re-introduction of coppicing will help to inject natural diversity and thus resilience into the woodland in light of climate change and increased pests and diseases. Planting will also be used on sites where regeneration does not meet an average of 3,000 stems/ha by year 10.

### **National Vegetation Classification**

















### Legend



© Crown copyright and database right [2016] Ordnance Survey [100021242]

> **Clearfell** coupes will simply be managed through clearcutting (of over 0.25ha) and restocked either through natural regeneration, replanting or a combination of the two.

**Coppicing** of hazel, oak and sweet chestnut stands will be used in discrete riparian areas. Coupes will be no larger than 0.5ha and will only be used where resources allow for adequate deer proof fencing. Standards will be retained where deemed as appropriate future crop trees.

**Minimum Interventions** are predominantly inaccessible or ecologically valuable areas where intervention will only occur to protect and ensure the future succession of key habitats and species.

**Open space** is managed to ensure forest cover does not exceed 2m in height, a tolerance of 20% forest cover will be accepted on some lower priority sites.

**Uniform shelterwoods** are predominately broadleaved domimated and ASNW sites which will be managed using seeding fellings with possible under planting of site suitable species to control light levels and develop good timber quality. **Irregular shelterwoods** will look to develop a complex CCF structure through the identification and thinning towards quality final crop trees for the future.

Single-tree selections are used on existing complex structured stands or sensitive sites often important for amenity value.

### **Broadleaf Thinning**

Broadleaf high forest will be assessed for thinning every 10 years with a visual inspection of the stand. Thinning will allow sub-dominant broadleaves sufficient light and space to mature or will release existing advanced regeneration. Younger patches of regeneration can be thinned to favour site native species with trees of good form and vigour being retained. Where broadleaves consist primarily of a single species, it may be possible to enlarge natural gaps through irregular thinning rather than create new gaps through group felling, however, in all cases the size of gap will be dependent on slope, aspect and site fertility and must not be detrimental to crop stability. These gaps will be utilised for enrichment planting using a mix of native species other than those occurring in the overstorey - rather than reliance on natural regeneration.

### **Conifer Thinning**

also help decide if an area of conifer is to be thinned or not. Light levels, existing ground vegetation and any evidence of natural regeneration also influences how many trees are marked for removal.

### Legend



Next thinning date 2016 2017 2018 2019 2020 2018 See page 15, Continuous Cover Forestry for more information

#### **Tamar Valley Forest Plan** 2016 - 2026 Page 15





### **Silviculture**

Areas of conifer are assessed for thinning every 5 years with the targeted removal of larch species a key objective. Other factors such as the quantity, condition, age and distribution of any broadleaf content will





PAWS managed under continuous cover (or shelterwood and selection) systems will be thinned to favour broadleaf components. The targeted removal of larch and western hemlock species will limit the impact of disease susceptibility to Phytophthora ramorum and the adverse impact dense w. hemlock has on natural regeneration. As a result this will increase the resilience of the woodlands as well as the regeneration potential of native species.

Uniform shelterwoods are predominately oak dominated sites which will be managed using seeding fellings, following the identification of final crop trees. Under planting of site suitable species, such as beech or hazel may be considered, all other mixed broadleaf stands will be managed irregularly through thinning. Irregular shelterwoods on PAWS will look to favour the development of regenerating native broadleaves and target the removal of mature conifer components.





© Crown copyright and database right [2016] Ordnance Survey [100021242]

#### **Tamar Valley Forest Plan** 2016 - 2026 Page 16





### **Continuous Cover Forestry**



### Legend

### Coupes

Open

| · · · · ·                   | Alternatives to Clearfell |
|-----------------------------|---------------------------|
|                             | Fell 2012 - 2016          |
|                             | Fell 2017 - 2021          |
|                             | Fell 2022 - 2026          |
|                             | Fell 2027 - 2031          |
|                             | Fell 2032 - 2036          |
|                             | Fell 2037 - 2041          |
|                             | Fell 2042 - 2046          |
|                             | Fell post 2046            |
| $\Diamond \Upsilon$         | Wood Pasture              |
| × * * *<br>× * * *<br>× * * | Retentions                |
|                             | Minimum Intervention      |
|                             | Natural Reserve           |



Coupe 83954 (0.51ha) Fell/Coppice 2022-26 (Broadleaves) Restock 82954a (0.51ha) 70% Broadleaves coppice D sam Denham 30% Retained broadleaf standards Hunter's Oak 6 ,

#### Restock 83749a (0.36ha) 70% Broadleaves coppice 30% Retained broadleaf standards

Fell/Coppice 2022-26 (Broadleaves)

Coupe 83749(0.36ha)

and the second second

AL Cole

Main Far

© Crown copyright and database right [2016] Ordnance Survey [100021242]

## **Management Prescriptions** 2016 - 2046

#### Tamar Valley Forest Plan 2016 - 2026 Page 17



PErc 















### Legend

Moor and Heath Hedgerow Forest Road Edge Conservation + Recreation Public Roadside •••• • Windthrow protection Watercourse Broadleaved Belt **Broad Habitat Type** BROADLEAVED; MIXED/YEW WOODLANDS **CONIFEROUS WOODLANDS Priority Habitat Type** Lowland mixed deciduous woodland





## **Conservation - Habitats**

Ride and road sides, together with watercourses and hedgerow management will conform to the prescriptions outlined in the District document, Design and Management of Environmental Corridors (Lucas, 2006). This document outlines the management of light levels, pinch points and forest edge dynamics.

The road and ride network within the Plan area will be utilised to extend and connect ride side habitats and transient open spaces, this will be achieved through targeted widening and unstocking of edges to some coupes following felling operations to create a mixed transient open and scrubby habitat for a multitude of species.

A variety of deadwood will be retained according to the level of ecological value and in line with Guidance (Humphrey & Bailey, 2012). Retaining decaying snags and logs as well senescent native trees throughout the woodland will create suitable deadwood habitat for numerous associated species.

The streamsides and wet woodland found at the bottom of valley remain predominantly wooded with either ASNW broadleaf woodland or planted Douglas fir. The majority of these sites will be managed through thinning and the recruitment of suitable wet woodland species such as alder, willow and birch encouraged. Where appropriate (notably Birch Wood) where secondary woodland sites are clearfelled, areas of open space and broadleaves will be included to add diversity and structure.

### Lowland Mixed Deciduous Woodland

A number of areas of remnant lowland mixed deciduous woodland (as shown right) are found across the Plan area. These are predominantly made up of Penduculate oak, ash, birch beech and sweet chestnut. Some evidence of coppicing of hazel and sweet chestnut exists.

Management of these areas will be sensitive to ensure the quality is maintained in perpetuity. Thinning will be reviewed on a ten yearly cycle with the aim of enhancing and improving the condition of the habitat. Removal of invasive or un-suitable species, such as laurel, rhododendron or Western hemlock will ensure that this habitat is maintained and used as a building block for future native broadleaf restoration.









**Trees of significance -** are found throughout the Plan area and will be retained for perpetuity. The majority of these trees are either oak (as shown) or sweet chestnut. When crops are thinned crowns will be released slowly to minimise the impact of sudden exposure to desiccating winds and sun scorch. Management will be in line with FC Guidance (Ops No. 31).



### **Conservation - Features**

The Forest Plan area is used by a vast array of common and rare flora and fauna. The considerable contribution the forests and their associated areas make to habitat provision in the landscape is widely recognised. Some flora and fauna species can have a detrimental impact on the forest and its features if their numbers are too high. Species such as rhododendron, wild deer and squirrel will all be managed in line with District Strategy to ensure that their pressure does not have a negative impact on the condition of habitats and crops.

**Otter** - are known to use the length of the River Tavy and is widespread across most rivers in Devon and Cornwall. This European Protected Species experienced a decline in previous decades but has recovered well in the south west of England. They inhabit streamside and wetland areas and the riparian woodland habitats found within the Plan area are ideal for nesting otter. The management of riparian wet woodland will be employed will ensure that a lush diversity of open space, scrub and high forest will ensure otter habitat is enhance and preserved to support this species.



**Raptor -** notably goshawk (above) and buzzard are known to nest and hunt within the forest areas. Many of the species choose to rest in high well branched conifer trees and then feed over open ground, making the forests ideal raptor habitat in an otherwise varied landscape. The management of appropriate large or potentially large trees for long retentions will ensure that habitat provision is maintained.



#### Tamar Valley Forest Plan 2016 - 2026 Page 20



onatry Communication woodlands have ween certified in Accordance with the view of the Forcet Newardship Council





**Dormouse** favourable habitat is found throughout the Plan area and therefore this species could be inhabiting the woodlands. This European Protected Species requires pinch points across corridors to allow habitat connectivity between broadleaved woodland, particularly in stands with a high hazel and/or sweet chestnut components. The increase in coppicing in hazel dominated stands will significantly enhance habitat quality with prescriptions outlined in the Environmental Corridors document also ensuring appropriate habitat provision and management will be in line with Best Practice Guidance (FC & NE,







Whilst no scheduled features are registered within the Plan area. The woodlands do play host to a significant amount of unscheduled features and cultural interest.



Wood.

These features are fenced but visible to the public maintaining their cultural value whilst keeping people safe. Mining areas are avoided during operations.



### Tamar Valley Forest Plan 2016 - 2026 Page 21





### **Heritage Features**

The copper mining industry made the Tamar Valley internationally famous in the 19th and early 20th century. A considerable number of copper mineshafts are found through the central spine of Birch Wood.

The mineshaft found in Denham was a silver and lead mine and pre-dates those of Birch





The Plan area experiences a small amount of low-key recreational usage. No Public Rights of Way cross the Plan area but Birch Wood is Open Access, with Denham being *de facto* Open Access due to the nature of the landholding.

The majority of recreation usage is made up of walkers with some limited amount of usage by horse riders and mountain bike riders.

The contribution the Plan area makes to the AONB as an area of recreation is significant and with facilities and trails maintained in to the future this value will be retained.



Birch Wood is held under freehold, having been acquired from Dartington Hall in 1957. As a result Open Access is provided in this part of the Plan area.

Denham is held under leasehold from Buckland Abbey Estate, also acquired in 1957, which is now owned by the National Trust. Access is limited to de facto.

The National Trust has retained the management of the Buckland Abbey Estate, notably the woodland to the east to Denham, Great North Wood. Management objectives and prescriptions will be in consultation and keeping with neighbouring woodland management.



### Legend

Denham\_CarPark

De Facto Access

**Restricted Byway** 

Class C Roads

**Open Access** 

Bridleway

---- Class A/B Roads

Byway

Rides

A Roads

**B** Roads

Minor Roads

Ρ

## **APPENDIX 1**



Denham 1 AL. Charles I Charles

© Crown copyright and database right [2016] Ordnance Survey [100021242]





Geology

Legend

HEAD

ALLUVIUM

**River Tavy** 

TAVY FORMATION - SLATE

## Tamar Valley Forest Plan 2016 - 2026 Page 23



## Vuð Esc





### Legend

- River Tavy
  - Unknown
  - 1 [Typical brown earth]
  - 1u [Upland brown earth]
  - 7 [Typical surface-water gley]



#### Tamar Valley Forest Plan 2016 - 2026 Page 24





## **Landform Analysis**

The landscape analysis is used to assess the landform patterns and demonstrates how it is in keeping with the surrounding landscape character.

One's eye is naturally dawn up the valleys and down the ridges. These principles will be used to design the shape of future coupes. Ensuring that the shape and size of felling and restocking areas do not detract from the natural appearance of the forest and its contribution to the landscape character,



## Landscape Analysis

Despite the minimal amount of planned clearfell within the Plan area an analysis of the implications of management proposals is required. This is to ensure the impact of the prescriptions does not adversely influence the local landscape. The integration of the woodlands and how they are managed with the local landscape is integral to the success of the Forest Plan and the retention of the landscape value of the Tamar Valley AONB.

The Forest Plan blocks can be seen from a number of viewpoints but is also often obscured or limited by the surrounding topography and features. Four key points of view have been identified for their exposure and impact and analysed below.

© Crown copyright and database right [2016] Ordnance Survey [100021242]



Improved agricultural fields intersect the woodland both in foreground and at a distance. This creates good integration between high forest and open pasture.









Areas of oak dominated ASNW sit at lower elevations. The shape of these intersects well with conifer with minimal geometric shapes or lines.

Planned clearfell for 2027-31 will have limited impact on the landscape due to the retention of surrounding crops.

Contiguous blocks of conifer are a key and prominent feature in the landscape with integration with broadleaf woodland, hedgerows and field system well established.









## **APPENDIX 2–Option Testing**

| <b>Option 1 – Current Forest Plan (Master)</b>  | <b>Option 2 – Proposed Forest Plan (Scenario)</b>   |
|---|---|
| The continued production of sustainable and marketable woodland products.   | I   |
| Significant peaks and troughs in production, mainly due to large clearfelling programme in 2027-31, potentially weaken the sustainability of the woodlands timber production. Continued production through thinning   | The felling and thinning programme is balanced across the deca<br>the sustainability of timber production.  |
| To protect enhance and restore areas of ancient woodland in line with the 2005  | Keepers of Time' policy.  |
| The Plan makes no reference to Keepers in Time, due to the time when it was written.<br>Management is through clearfell and restock with conifer. This is wholly incompatible with<br>the Policy.   | Plan makes significant and clear provision for the restoration of gradual thinning and enhancement of broadleaf remnants. This over time whilst retaining ecosystem functioning.                      |
| Deliver well-designed forests in keeping with the AONB and local landscape chara  | acter.  |
| Whilst the original clearfell programme was well integrated into the landscape, odd clearfell coupe shapes following multiple amendments threaten the value of the local area. The continued use of Douglas fir restock contributes to the AONB asset but fails to meet desire and policy to restore to native broadleaf cover. | The move away from clearfell is in keeping with the local landsc<br>acknowledgement of the value of Douglas fir to the local charac<br>gradual reduction is in line with local aspiration and Policy. |
| Protect and enhance woodland and open habitats and their associated species.  |   |
| The Plan makes provision for open space and other valuable habitats such as ASNW but does not make them a core objective  | The Plan identifies and prescribes management for current and and corridors.  |
| To conserve, maintain and enhance cultural and heritage assets.   | ·   |
| Plan makes no mention or provision for the protection and enhancement of any heritage assets, unscheduled or unknown.   | The small number of heritage features, mineshafts, are identified consideration in the planning of future management.   |
| The provision and maintenance of recreation facilities.   |   |
| Plan makes minimal reference or significance of recreational facilities.  | The Plan acknowledges the value of the woodland to low key re-<br>takes steps to ensure this is maintained.   |





#### **Tamar Valley Forest Plan** 2016 - 2026 Page 27

rsc FSC















#### Name: Phytophthora ramorum (PR)

First appearance: 2009

#### Attacks: Larches

P. ramorum was first found in the UK in 2002 and until 2009 in the woodland environment had largely been associated with rhododendron species acting as a host from which spores are produced. In August 2009 P. ramorum was found on a small number of dead and dving Japanese Larch in South West England, causing particular concern since some affected trees were not close to infected rhododendron and showing a significant change in the dynamics of the disease than experienced previously. Following this testing in Devon and west Somerset confirmed the presence of PR in mature Japanese larch as well as species in its under-storey, including sweet chestnut, beech, birch, oak, Douglas fir and Western hemlock. On some sites there is little or no rhododendron present. It is now known that Japanese larch can produce very high quantities of diseasecarrying spores when actively growing in spring and sum-



© Crown copyright and database right [2016] Ordnance Survey [100021242]

mer, at much higher levels than those produced by rhododendron. These can be spread significant distances in moist air. PR is a notifiable disease dealt with by felling the infected area under a statutory plant health notice (SPHN) issued through FERA and the Forestry Commission.





#### Name: Oak 'dieback' or 'decline'

First appearance: unknown

#### Affects: Oak

Oak 'dieback' or 'decline' is the name used to describe poor health in oak trees and can be split into Chronic decline and Acute decline. Chronic decline is protracted taking effect on the Oak over a number of decades whilst Acute decline is much swifter acting over much shorter periods usually five years or so. Symptoms can be caused by a range of living agents e.g. insect and fungal attack, or non-living factors, e.g. poor soil and drought. Factors causing decline can vary between sites, as can the effects of the factors through time. Oak decline is not new; oak trees in Britain have been affected for the most part of the past century. Both native species of oak are affected, but Pedunculate oak (Quercus robur) more so than Sessile oak (Quercus petraea). Successive exposure to any of these agents on a yearly/seasonal basis further reduces the health of the tree(s) and predisposes it to other living (Biotic) agents that can often spell the eventual death knell for the tree.





### Name: Dothistroma Needle Blight (DBN)

First appearance: mid 1990s

Attacks: Pine species

Often referred to as Red Band Needle Blight (RBN) and can reduce growth rates by between 70 and 90%. Effects of RBN are managed through thinning the wood more heavily than you would normally to introduce higher levels of air flow through the remaining crop. However, the Mortimer Plan area contains a relatively small component and therefore its impact has been fairly limited.





| Term                                   | Abbreviation          | Description  |
|--|-----------------------|--|
| Ancient Semi-<br>Natural Wood-<br>land | ASNW                  | An ancient woodland site, where trees and other plant species appear to of established naturally rather than having been nantly these sites will contain 80% or over of site native species or species native to the surrounding area.   |
| Alternatives to<br>Clearfell           | ATC                   | Alternative to Clearfell is similar to CCF and refers to management systems where stands are regenerated without clearfe   |
| Ancient Wood-<br>land Site             | AWS                   | A site that has technically been wooded since 1600AD and is unlikely to have been converted to farmland in the last few  |
| Continuous<br>Cover Forestry           | CCF                   | Continuous Cover Forestry is an approach to forest management that enables an owner of woodland to manage the wood<br>need for clearfelling. This enables tree cover to be maintained, usually with one or more levels and can be applied to bot<br>stands. With Conifer it is possible to regenerate the crop a lot faster than in broadleaf crops, where the canopy is genera<br>slower and over a much longer time span. A decision to use CCF must be driven by management objectives and will have<br>ten aimed at creating a more diverse forest, both structurally and in terms of species composition. There are no standard<br>ing CCF is very flexible in ensuring opportunities can be taken advantage of as they arise. This development of a more diverse<br>sible way to reduce the risks posed by future changes in the climate and biotic threats.   |
| Clearfell                              | C/F or CF             | To cut and remove all trees from a certain area of woodland.   |
|  |                       | A stand of trees. Often associated with stands completely or partially managed for its timber.   |
| Сгор                                   |                       | Just as farmers manage crops so does forestry the only difference is a farmers' rotation is shorter and often realised in 1 much longer term crop with rotations varying from 6 years to 400 years. (also see definition for rotation)   |
| Enrichment<br>planting                 |                       | Planting different species within areas of regen that helps diversify the range of species in a wood and in doing so can ma<br>future climate change and future threats from disease.<br>Enrichment may be desirable in areas where success of regeneration is uneven, patchy or where a regen crop is limited b<br>cies present.  |
| Group felling /<br>group planting      |                       | This is where small areas of woodland are felled hence the name "group felling" and then either allowed to develop throug<br>regen or in this case planted hence "group planting". These techniques can help to develop structure* within a wood ove<br>time and is often used in conjunction with continuous cover. *Either in terms of age or number of tree species present,<br>shade are provided by the remaining upper storey one can consider a larger number of tree species when deciding what t  |
| Hectare                                | На                    | Unit of area equating to 2.47 acres.   |
| Native (and<br>honorary na-<br>tive)   |                       | The trees making up the woodland are part of England's natural, or naturalised flora. Determined by whether the trees of out assistance from humans since the last ice age (or in the case of 'honorary natives' were brought here by people but h historic times); and whether they would naturally be found in this part of England.   |
| Natural Regen-<br>eration              | Regen or<br>nat-regen | Trees growing on a site as a result of natural seed fall, and can be used as a management process and can allow cleared germinate, grow and develop naturally. This process can happen anywhere and woods can be managed to encourage na there is no guarantee of success. In these instances, or if nat-regen is unlikely for a variety of reasons, one can use enrice group planting to achieve the same affect. The process usually relies on an overstorey of "parent trees" being present or on parent trees being close by to provide t ent trees will usually of been thinned and managed with natural regeneration in mind. Existing areas of nat-regen are then usually developed through carefully thinning the surrounding woodland over a number of 'recruited') into the main crop for the next rotation at some point in the future. Usually done in small groups or in strips this system can allow a varied woodland structure to develop over time. Protection from competing plant species and mammal browsing might be required in the early stages by fencing or using |

planted. Predomi-

### elling.

centuries.

dland without the th conifer or broadleaf ally removed a lot ve long-term vision ofrd prescriptions meanliverse forest is a sen-

year. Trees are a

ake it more resilient to

by the number of spe-

igh the use of nater a given length of since shelter and to plant.

colonised Britain withhave naturalised in

areas of woodland to at-regen although ichment planting or

the seed. These par-

ber of years, to give incorporated

tree shelters.

#### Tamar Valley Forest Plan 2016 - 2026 Page 30



woodlânde have neen certitiert in accordance with the rules of the Forest Stewardship Council



## APPENDIX 3— Glossary

|              |    | Generally a commercial term used to describe the length of time an area of trees is growing for, from the time of planting<br>For broadleaves a rotation is generally a lot longer than that of conifer species* and can broadly speaking be anywhere b<br>years, as opposed to conifer crops whose rotation is generally shorter but can vary from 20-25 years to 120 years plus.  |
|--------------|----|---|
| Rotation     |    | *The exception being that of coppice where rotation length can vary from 5 or 6 years up to 30 years plus depending on  |
|              |    | "First rotation" would refer to an area of wood planted on open ground not previously wooded. And so "second rotation" is has been cleared and replanted.   |
| Shelterwood  |    | A management system that is applicable to conifer or broadleaf, where tree canopy is maintained at one or more levels w<br>fell the whole site. Felling can occur, but generally in small "groups" whose size shape and spatial distribution will vary de<br>tions. The "groups" are then either: allowed to develop and establish by the use of natural regeneration, are planted or a<br>mixture of both techniques. This known as a "group shelterwood system"   |
|              |    | A variation on this is "Single tree selection". This variation removes individual trees of all size classes more or less uniform<br>to maintain an uneven-aged stand and achieve other stand structural objectives. While it is easier to apply such a system<br>rally close to the uneven-aged condition, single tree selection systems can be prescribed for even-aged stands, although<br>thinning interventions must be made to create a stand structure where the system can truly be applied.   |
| Silviculture |    | A term coined during late 19th century from the Latin <i>silva meaning</i> 'wood' and the French <i>culture</i> meaning 'cultivation' a art and science of controlling the establishment, growth, composition, and quality of forest vegetation to achieve a full rai jectives.   |
| Stand        |    | A group or area of trees that are more or less homogeneous with regard to species composition, density, size, and someti  |
| Thin         | ТН | <ul> <li>Selective removal of trees from a wooded area, giving remaining trees more space to grow into larger trees. Thinning is a Improve the quality and vigour of remaining trees.</li> <li>Remove trees interfering with mature or veteran broadleaf trees.</li> <li>Give space for tops (or "crowns") of broadleaf trees to develop and potentially act as a future seed source.</li> <li>Give space for natural regeneration to grow and develop with the intention of recruiting these younger naturally grown tree ture woodland structure.</li> <li>Create gaps for group planting or enrichment.</li> <li>Remove species of tree that may compromise the intended management objective of the woodland eg: non-native or invasionation system.</li> <li>Improve the economic value of a wood.</li> <li>Help realise opportunities to enhance ecological value.</li> </ul> |
| Yield Class  | YC | A method of measuring the growth rate or "increment" of a crop of trees by age and height; measured in m3 per Ha per a YC of 16 is one that has an annual increment of more than 16m3 but less than 17m3, although generally only even nun stating YC.  |

#### Tamar Valley Forest Plan 2016 - 2026 Page 31

g to the time of felling. Detween 80 years to 3-400

management objectives.

is one where woodland

without the need to clearlepending on site condiare established using a

rmly throughout the stand n to a stand that is natunumerous preparatory

and so Silviculture is the ange of forest resource ob-

imes habitat.

done to:

rees as a part of the fu-

vasive species such as

annum. E.g. A crop with mbers are used when

| The sector of |
|---------------|

Foreatry Commission woodlends have been contribut in occur during with the rules of the Forest Stewardship Council



### References

Forestry Commission, 2011, The UK Forestry Standard, Forestry Commission, Edinburgh

Forestry Commission, 2005, Keepers in Time: A Statement of Policy for Englands Ancient and Native Woodlands, Forestry Commission, Cambridge Forestry Commission & Natural England, 2007, Guidance on managing woodlands with dormice in England, Forestry Commission, Bristol Forestry Commission, 2013a, West England Forestry District Strategy 2013-2020, Forestry Commission, Bristol Forestry Commission, 2013b, Strategic Plan for the Public Forest Estate in England, Forestry Commission, Bristol Humphrey, J. & Bailey, S., 2012, *Managing deadwood in forests and woodlands*, Forestry Commission, Edinburgh Lucas, O., 2006, Design and Management of Environmental Corridors, Peninsula Forest District, Forestry Commission, Exeter Natural England, 2014, National Character Assessment, 151 South Devon Natural England, York Tamar Valley AONB, 2014, AONB Management Plan, AONB, Plymouth UKWAS, 2012, United Kingdom Woodland Assurance Standard, UKWAS, Edinburgh West Devon District Council (2008) *Character Description — River Tavy Middle Valley*, West Devon District Council, Tavistock



